

# Small Odd Cycles in 4-Chromatic Graphs

---

**Tao Jiang**

DEPARTMENT OF MATHEMATICAL SCIENCES  
MICHIGAN TECHNOLOGICAL UNIVERSITY  
HOUGHTON, MICHIGAN 49931  
E-mail: [tjiang@mtu.edu](mailto:tjiang@mtu.edu)

Received August 3, 1999

**Abstract:** It is shown that every 4-chromatic graph on  $n$  vertices contains an odd cycle of length less than  $2\sqrt{n}+3$ . This improves the previous bound given by Nilli [J Graph Theory 3 (1999), 145–147]. © 2001 John Wiley & Sons, Inc.  
J Graph Theory 37: 115–117, 2001

Keywords: *odd cycle; chromatic number; girth*

By a theorem of Kierstead, Szemerédi, and Trotter [1], every 4-chromatic graph on  $n$  vertices contains an odd cycle of length at most  $8\sqrt{n}$ . On the other hand, there are infinitely many values of  $n$  for which there are 4-chromatic graphs on  $n$  vertices containing no odd cycle of length less than  $\sqrt{2n}$ . Constructions of such graphs appear in [4,5,2], where, in the last article, the authors raise the natural problem of determining the supremum of the reals  $c$  such that there are infinitely many values of  $n$  for which there are 4-chromatic graphs on  $n$  vertices with no odd cycles of length less than  $c\sqrt{n}$ . By the above results, it is known that this supremum is between  $\sqrt{2}$  and 8. Recently, Nilli [3] improved the upper bound by showing that every 4-chromatic graph on  $n$  vertices contains an odd cycle of length less than  $\sqrt{8n}$ . Hence, the supremum is now between  $\sqrt{2}$  and  $\sqrt{8}$ . In this

---

1990 Mathematics Subject Classification 05C35, 05C55

© 2001 John Wiley & Sons, Inc.

note, we further improve the upper bound to 2. Our method is based on the one Nilli uses.

**Theorem 1.** *Every 4-chromatic graph on  $n$  vertices contains an odd cycle of length less than  $2\sqrt{n} + 3$ .*

**Proof.** Let  $G$  be a 4-chromatic graph on  $n$  vertices. Suppose the shortest odd cycle of  $G$  has length  $2k + 3$ , we prove that  $n > k^2$ , from which we have  $k < \sqrt{n}$ , and the theorem follows.

Let  $A$  and  $B$  be two disjoint independent sets in  $G$  for which the cardinality  $|A \cup B|$  is maximum. Since  $G$  is 4-chromatic,  $G - A$  contains an odd cycle. Let  $C$  be a shortest odd cycle in  $G - A$ ; clearly it is an induced cycle. By our assumption,  $C$  has length at least  $2k + 3$ . Let  $v$  be a vertex on  $C$ . For each integer  $i \geq 0$ , define  $V_i = \{u \in V(G) : d(u, v) = i\}$ , where  $d(u, v)$  is the distance between  $u$  and  $v$  in  $G$ . In particular,  $V_0 = \{v\}$ . Since  $G$  has no odd cycle of length at most  $2k + 1$ ,  $V_i$  is an independent set for each  $i \leq k$ . For each  $i$ ,  $1 \leq i \leq k$ , define  $H_i$  to be the subgraph of  $G$  induced by

$$U_i = (\cup_{0 \leq j < i} V_j) \cup (V_i \cap B) \cup (\cup_{j > i} (V_j \cap (A \cup B))) \tag{1}$$

We claim that  $H_i$  is bipartite. Let  $S = (\cup_{0 \leq j < i} V_j)$  and  $T = \cup_{j > i} (V_j \cap (A \cup B))$ . Note that there is no edge between  $S$  and  $T$ , since vertices in  $S$  are at distance at most  $i - 1$  from  $v$ , whereas vertices in  $T$  are at distance at least  $i + 1$  from  $v$ . Since  $V_j$  is an independent set for each  $j \leq i$ , the subgraph  $H'_i$  induced by  $S \cup (V_i \cap B)$  is bipartite with a bipartition  $X_1, Y_1$ , such that  $(V_i \cap B) \subseteq Y_1$ . On the other hand, the subgraph induced by  $T$  is also bipartite with a bipartition  $X_2 = T \cap A, Y_2 = T \cap B$ . Since there is no edge between  $S$  and  $T$ , and  $B$  is an independent set, the only possible edges between  $H'_i$  and  $H''_i$  are between  $V_i \cap B$  and  $T \cap A = X_2$ . Since  $V_i \cap B \subseteq Y_1$  by our assumption,  $H_i$  is bipartite with a bipartition  $X_1 \cup X_2, Y_1 \cup Y_2$ . It follows that  $U_i$  is the union of two disjoint independent sets. By our choice of  $A, B$ , we have  $|A \cup B| \geq |U_i|$ . In particular,  $|A \cup B - U_i| \geq |U_i - A \cup B|$ . Note that  $U$  contains  $2i - 1$  vertices on  $C$  that are within distance  $i - 1$  from  $v$ . These vertices induce a path in  $G - A$ . Since  $B$  is an independent set, it contains at most  $i$  of those vertices, hence at least  $i - 1$  of those vertices do not belong to  $A \cup B$ . This shows that  $|U_i - A \cup B| \geq i - 1$ . On the other hand, it is easy to see that  $A \cup B - U_i = V_i \cap A$ . It follows that  $|V_i \cap A| = |A \cup B - U_i| \geq |U_i - A \cup B| \geq i - 1$ . Hence, we have

$$|A| \geq \sum_{i=1}^k |V_i \cap A| \geq \sum_{i=1}^k (i - 1) = k(k - 1)/2 \tag{2}$$

By symmetry, we have  $|B| \geq k(k - 1)/2$ . Observe also that at least  $k + 2$  vertices on  $C$  do not lie in  $B$ , and hence not in  $A \cup B$ . Therefore,

$$n \geq |A| + |B| + (k + 2) = k^2 + 2 > k^2 \tag{3}$$

This completes our proof. ■

**References**

- [1] H. Kierstead, E. Szemerédi, and W. T. Trotter, On coloring graphs with locally small chromatic number, *Combinatorica* 4 (1984), 183–185.
- [2] N. V. Ngoc and Zs. Tuza, 4-chromatic graphs with large odd girth, *Discrete Math* 138 (1995), 387–392.
- [3] A. Nilli, Short odd cycles in 4-chromatic graphs, *J Graph Theory* 31 (1999), 145–147.
- [4] H. Sachs and M. Stiebitz, Constructive methods in the theory of color-critical graphs, *Discrete Math* 74 (1989), 201–226.
- [5] D. A. Youngs, 4-chromatic projective graphs, *J Graph Theory* 21 (1996), 219–227.